

Dirac Equation in External Non-Abelian Gauge Field in Plane Wave Approximation

A.V.Koshelkin.*

Moscow Institute for Physics and Engineering, Kashirskoye sh., 31, 115409 Moscow, Russia

(Dated: July 19, 2009)

The exact solutions of the Dirac equation in an external non-abelian $SU(N)$ gauge field which is in the form of a plane wave on the light cone is obtained. The whole set of the solutions for both particles and anti-particle is constructed.

PACS numbers: 11.15.-q, 11.10.-z, 03.70.+k

I. INTRODUCTION

The study of non-abelian gauge fields plays an important role in the modern field theory[1, 2, 3]. This primarily concerns the field governed by unitary symmetries. Exactly such symmetry ($SU(N)$) is the basis of QCD[4] where quarks form the space of the fundamental representation of $SU_c(3)$ group while gauge fields provide interaction between them. Therefore the knowledge of the solution of the Dirac equations in such field is the key to understanding the processes taking place in the strong interacting matter appearing, particular, in the collisions of heavy ions of high energies[5].

The Dirac equation in the field of a plane electromagnetic field wave has been studied by D.M.Volkov who has derived the exact solution[6] for the first time. The key point in obtaining the such solution is the commutativity of electromagnetic field.

In the present paper the exact solution of the Dirac equation in an external non-abelian $SU(N)$ gauge field which represents a plane wave on the light cone is obtained. The whole set of the solutions which determines the states of both particles and anti-particles is derived. On a basis of the obtained solutions the fermion current is calculated. It strongly differs in structure from the fermion current in the field of a plane wave in QED[6].

The paper is organized as follows. The statement of the considered problem is formulated in Section II. Section III contains the solution of the Dirac equation in an external non-abelian gauge field. The last section is conclusion. The transformation of the operator exponent is presented in Appendix.

II. STATEMENT OF THE PROBLEM

We consider a fermion with a spin $s = 1/2$ in an external non-abelian field A_a^ν . The fermion fields $\Psi(x)$ and field A_a^ν form the space of the fundamental ($\Psi(x)$) and associated (A_a^ν) representations of $SU(N)$ group, respectively. Then the fermion field $\Psi(x)$ satisfies the equations[7, 8]:

$$\{i\gamma^\mu (\partial_\mu + ig \cdot A_\mu^a(x)T_a) - m\} \Psi(x) = 0 \quad (1)$$

$$\bar{\Psi}(x) \left\{ i\gamma^\mu \left(\overleftarrow{\partial}_\mu - ig \cdot A_\mu^a(x)T_a \right) + m \right\} = 0; \quad \bar{\Psi}(x) = \Psi^\dagger(x)\gamma^0, \quad (2)$$

where m is a fermion mass, g is the coupling constant; γ^ν are the Dirac matrixes, $x^\mu = (x^0; \vec{x})$ is a vector in the Minkowski space-time; $\partial_\mu = (\partial/\partial t; \nabla)$; the Roman letters numerate the basis in the space of the associated representation of $SU(N)$ group, so that $a, b, c = 1 \dots N^2 - 1$. We use the signature $diag(\mathcal{G}^{\mu\nu}) = (1; -1; -1; -1)$ for the metric tensor $\mathcal{G}^{\mu\nu}$. The line over Ψ and "dagger" mean the Dirac and hermitian conjugation, respectively[6, 7]. Summing over any pair of repeated indexes is implied.

The symbols T_a in Eqs.(1), (2) are the generators of $SU(N)$ group which satisfy the commutative relations and normalization condition:

*koshelkin@mtu-net.ru; koshelkin@theor.mephi.ru;

$$[T_a, T_b]_- = T_a T_b - T_b T_a = i f_{ab}^c T_c; \quad (3)$$

$$Tr (T_a T_b) = \frac{1}{2} \delta_{ab}; \quad (4)$$

where f_{ab}^c are the structure constant of the $SU(N)$ group, which are real and anti-symmetrical with respect to the transposition in any pair of indexes; δ_{ab} is the Kroneker symbol. In the matrix representation the operators T_a coincide with the Pauli and Gill-Mann matrixes when N is equal to 2 or 3, respectively

It directly follows from Eqs.(3), (4) that

$$[T_a, T_b]_+ = T_a T_b + T_b T_a = \frac{N^2 - 2}{2N} \delta_{ab}. \quad (5)$$

Let the fields $A_\mu^a(x)$ be dependent on the 4-coordinate x in the form of a plane wave:

$$A_\mu^a(x) = A_\mu^a(kx) \equiv A_\mu^a(\varphi); \quad k^\mu k_\mu = 0; \quad \varphi = k_\mu x^\mu \equiv kx, \quad (6)$$

where $k = (k^0; \vec{k})$ is some 4-vector lying on the light cone.

We fix the axial gauge by the expressions:

$$\partial^\mu A_\mu^a = 0; \quad k^\mu \dot{A}_\mu^a = 0, \quad (7)$$

where the dot over the letter means differentiation with respect to the variable $\varphi = kx$.

III. SOLUTION OF DIRAC EQUATION IN EXTERNAL NON-ABELIAN GAUGE FIELD

To derive the fermion field $\Psi(x)$ we go from Eq.(1) to the so-called quadrated Dirac equation which has the following form:

$$\begin{aligned} & \left\{ \partial_\mu \partial^\mu - m^2 + g^2 (\gamma^\mu A_\mu^a T_a)^2 - 2ig (\gamma^\mu A_\mu^a T_a) (\gamma^\mu \partial_\mu) - ig (\gamma^\mu k_\mu) (\gamma^\mu \dot{A}_\mu^a T_a) \right\} \Phi(x) = 0; \\ \Psi(x) &= \left\{ \frac{i\gamma^\mu (\partial_\mu + ig \cdot A_\mu^a(x) T_a) + m}{2m} \right\} \Phi(x). \end{aligned} \quad (8)$$

We will find the solution of the last equation in the form:

$$\Phi(x) \equiv \Phi_{\sigma,\alpha}(x, p) = e^{-ipx} \cdot F_{\sigma,\alpha}(kx). \quad (9)$$

where $F_{\sigma,\alpha}(kx)$ is some multicomponent function which is a generated Dirac spinor. It depends on both the spin variable σ and the variable α which specifies the state of a fermion in the space of the fundamental representation of the $SU(N)$ group, thereat $\alpha = 1 \div N$. $p^\nu = (p^0, \vec{p})$ is the 4-momentum of a particle.

We substitute $\Phi_{\sigma,\alpha}(x, p)$ in the form given by Eq.(9) into the formula (8). Using the relations for the γ - matrixes[6] and Eq.(5), we obtain:

$$\begin{aligned} & \left\{ p^2 - m^2 + g^2 \frac{N^2 - 2}{4N} (A_\mu^a A_\mu^a) - 2g (T_a A_\mu^a p^\mu) - ig (\gamma^\mu k_\mu) (\gamma^\mu T_a \dot{A}_\mu^a) \right\} F_{\sigma,\alpha}(\varphi) + 2i(pk) \dot{F}_{\sigma,\alpha}(\varphi) = 0; \\ \varphi &= kx, \quad (pk) = p^\mu k_\mu, \end{aligned} \quad (10)$$

where the dot over $\dot{F}_{\sigma,\alpha}(\varphi)$ means derivative with respect to the variable φ as before.

The solution of the derived equation can be formally written as follows:

$$F_{\sigma,\alpha}(\varphi) = \frac{1}{\sqrt{2p^0}} \exp \left\{ i \frac{(p^2 - m^2)\varphi + g^2 \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a(\varphi') A_\mu^a(\varphi'))}{2(pk)} \right\} \\ \exp \left\{ -ig T_a \frac{\int_0^\varphi d\varphi' (A_\mu^a p^\mu) + \frac{i}{2} (\gamma^\mu k_\mu) (\gamma^\mu A_\mu^a)}{(pk)} \right\} u_\sigma(p) \cdot v_\alpha, \quad (11)$$

where $u_\sigma(p)$ and v_α are some spinors which are the elements of the spaces of the corresponding representations.

The last exponent in Eq.(11) is the operator. It is necessary to determine acting such operator on the spinors $u_\sigma(p)$ and v_α . To do this we expanse the second exponent in Eq.(11) in the series. After that, we group the odd and even terms separately. We also note, that due to Eqs.(6), (7) there are not any term containing γ -matrix in the power more than the first one in the series. The commutative relation (5) as well as Eqs.(6), (7) allow us to reduce the series which correspond to the odd and even terms in the expansion of the exponent to the combinations of sinuses and cosines. As a result we derive:

$$\exp \left\{ -ig T_a \frac{\int_0^\varphi d\varphi' (A_\mu^a p^\mu) + \frac{i}{2} (\gamma^\mu k_\mu) (\gamma^\mu A_\mu^a)}{(pk)} \right\} = \\ \cos \theta \left\{ \left(1 - ig T_a \frac{\tan \theta}{\theta(pk)} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \right) + \frac{g (\gamma^\mu k_\mu) (\gamma^\mu A_\mu^a)}{2(pk)} \cdot \left[\frac{\tan \theta}{\theta} T_a + \right. \right. \\ \left. \left. \frac{g}{(pk)} \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \left(-i \frac{\tan \theta}{\theta} + \frac{g}{(pk)} \frac{\theta - \tan \theta}{\theta^3} T_b \int_0^\varphi d\varphi' (A_\mu^b p^\mu) \right) \right] \right\}; \\ \theta = \frac{g}{(pk)} \sqrt{\frac{N^2-2}{4N}} \left(\int_0^\varphi d\varphi' (A_\mu^a(\varphi') p^\mu) \int_0^\varphi d\varphi' (A_\mu^a(\varphi'') p_\mu) \right)^{\frac{1}{2}} \quad (12)$$

Substituting the exponent given by the last formula into Eq.(11) we obtain:

$$\Phi_{\sigma,\alpha}(x, p) = e^{-ipx} F_{\sigma,\alpha}(\varphi) = e^{-ipx} \cdot \frac{1}{\sqrt{2p^0}} \exp \left\{ i \frac{(p^2 - m^2)\varphi + g^2 \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a(\varphi') A_\mu^a(\varphi'))}{2(pk)} \right\} \\ \cos \theta \left\{ \left(1 - ig T_a \frac{\tan \theta}{\theta(pk)} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \right) + \frac{g (\gamma^\mu k_\mu) (\gamma^\mu A_\mu^a)}{2(pk)} \cdot \left[\frac{\tan \theta}{\theta} T_a + \right. \right. \\ \left. \left. \frac{g}{(pk)} \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \left(-i \frac{\tan \theta}{\theta} + \frac{g}{(pk)} \frac{\theta - \tan \theta}{\theta^3} T_b \int_0^\varphi d\varphi' (A_\mu^b p^\mu) \right) \right] \right\} u_\sigma(p) \cdot v_\alpha, \quad (13)$$

To determine the spinor $u_\sigma(p)$ we assume that at time moment $t = -\infty$ there is no interaction of the fermion with the external field. Then, $u_\sigma(p)$ is the standard Dirac spinor satisfying the free Dirac equation. Thereat, we should note that the fermion is on-shell, so that $p^2 = m^2$. Since the spinor $u_\sigma(p)$ is independent on the time variable it is defined as the spinor of the free Dirac field[6] at any time moment.

We take the following normalization for $u_\sigma(p)$:

$$\bar{u}_\sigma(p) u_\lambda(p') = \pm 2m \delta_{\sigma\lambda} \delta_{pp'}; \quad p^2 = m^2, \quad (14)$$

where the plus and minus signs correspond to the Dirac scalar production of the spinors $u_\sigma(p)$ and $u_\sigma(-p)$, respectively. As for the spinor v_α we determine it by the relations:

$$v_\alpha^\dagger v_\beta = \delta_{\alpha\beta}; \quad Tr(T_a) = 0; \quad Tr(T_a T_b) = \frac{1}{2}\delta_{ab} \quad (15)$$

Substituting Eq.(13) into the formula (8) we derive the following after the direct calculations:

$$\begin{aligned} \Psi(x) = \Phi_{\sigma,\alpha}(x, p) &= \frac{e^{-ipx}}{\sqrt{2p^0}} \exp \left\{ i \frac{g^2 \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a(\varphi') A_\mu^a(\varphi'))}{2(pk)} \right\} \\ &\cos \theta \left\{ \left(1 - ig T_a \frac{\tan \theta}{\theta(pk)} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \right) + \frac{g (\gamma^\mu k_\mu) (\gamma^\mu A_\mu^a)}{2(pk)} \cdot \left[\frac{\tan \theta}{\theta} T_a + \right. \right. \\ &\left. \left. \frac{g}{(pk)} \frac{N^2-2}{4N} \int_0^\varphi d\varphi' (A_\mu^a p^\mu) \left(-i \frac{\tan \theta}{\theta} + \frac{g}{(pk)} \frac{\theta - \tan \theta}{\theta^3} T_b \int_0^\varphi d\varphi' (A_\mu^b p^\mu) \right) \right] \right\} u_\sigma(p) \cdot v_\alpha; \\ \theta &= \frac{g}{(pk)} \sqrt{\frac{N^2-2}{4N}} \left(\int_0^\varphi d\varphi' (A_\mu^a(\varphi') p^\mu) \int_0^\varphi d\varphi'' (A_\mu^a(\varphi'') p_\mu) \right)^{\frac{1}{2}}; \quad p^2 = p^\mu p_\mu = m^2. \end{aligned} \quad (16)$$

The derived $\Psi(x)$ -function is the exact solution of the Dirac equation in the external non-abelian field having the form of a plane wave on the light cone. The function (16) is normalized by the δ -function as follows:

$$\int d^3x \Psi^\dagger(x, p') \Psi(x, p) = (2\pi)^3 \delta^3(\vec{p} - \vec{p}'). \quad (17)$$

The direct calculation shows that $\Phi_{\sigma,\alpha}(x, p)$ and $\Phi_{-\sigma,\alpha}(x, -p)$ are orthogonal. In this way, it is obvious, that $\Phi_{\sigma,\alpha}(x, p)$ is the so-called positive frequency function whereas $\Phi_{-\sigma,\alpha}(x, p)$ is negative frequency one[6, 7]. This fact allows us to construct the general solution of the Dirac equation which describes the states both particles and anti-particles. Combining the functions $\Phi_{\sigma,\alpha}(x, p)$ and $\Phi_{-\sigma,\alpha}(x, -p)$, we obtain:

$$\begin{aligned} \Psi(x) &= \sum_{\sigma,\alpha} \int \frac{d^3p}{\sqrt{2p^0} (2\pi)^3} \left\{ \hat{a}_{\sigma,\alpha}(p) \Psi_{\sigma,\alpha}(x, p) + \hat{b}_{\sigma,\alpha}^\dagger(p) \Psi_{-\sigma,\alpha}(x, -p) \right\} \\ \bar{\Psi}(x) &= \sum_{\sigma,\alpha} \int \frac{d^3p}{\sqrt{2p^0} (2\pi)^3} \left\{ \hat{a}_{\sigma,\alpha}^\dagger(p) \bar{\Psi}_{\sigma,\alpha}(x, p) + \hat{b}_{\sigma,\alpha}(p) \bar{\Psi}_{-\sigma,\alpha}(x, -p) \right\}, \end{aligned} \quad (18)$$

where the symbols $\hat{a}_{\sigma,\alpha}^\dagger(p); \hat{b}_{\sigma,\alpha}^\dagger(p)$ and $\hat{a}_{\sigma,\alpha}(p); \hat{b}_{\sigma,\alpha}(p)$ are the operators of creation and cancellation of a fermion ($\hat{a}_{\sigma,\alpha}(p); \hat{a}_{\sigma,\alpha}^\dagger(p)$) and anti-fermion ($\hat{b}_{\sigma,\alpha}(p); \hat{b}_{\sigma,\alpha}^\dagger(p)$), respectively[6, 7]. Thereat, $\hat{a}_{\sigma,\alpha}(p) \hat{a}_{\sigma,\alpha}^\dagger(p); \hat{b}_{\sigma,\alpha}(p) \hat{b}_{\sigma,\alpha}^\dagger(p)$ satisfy the standard commutative relations for fermion operators.

IV. CONCLUSION

A relativistic fermion with the spin 1/2 in a external $SU(N)$ gauge field is considered. The exact solution of the Dirac equation where the field is in the form of a plane wave on the light cone is derived. The whole set of the solution which corresponds to both particles and anti-particles is obtained.

- [2] D.Gross, Chinese J. of Physics, **30**, 955 (1992).
- [3] A.M.Polyakov, Gauge Field and Strings. Harwood Academic Publishers, Chur, Switzerland, 1987
- [4] F.J.Yndurain. The Theory of Quarks and Gluons Interaction, Springer, 2006.
- [5] D.E.Kharzeev, Nucl.Phys. **A 699** 95 (2002); D.E.Kharzeev, J.Raufeisen, nucl-th/0206073, v1, 28 June, 2002.
- [6] V.B.Berestetskii, E.M.Lifshitz, L.P.Pitaevskii. Quantum Electrodynamics, Pergamon Press, 1981.
- [7] C.Itzykson, J.-B.Zuber. Quantum Field Theory. McGraw-Hill Book Company, 1980.
- [8] A.Slavnov, L.Faddeev, Introduction to Quantum Theory of Gauge Fields, 2nd enl. and rev. ed. Moscow, Nauka, 1988.